




# PASTE BAIT RODENTICIDE

**3kg** contains 15g sachets

ACTIVE INGREDIENT:  
Brodifacoum .....0.05g/kg

Registered by / Geregistreer deur:  
Innovative Pest Management  
Co. Reg. No. /Mpy. Reg. Nr.:  
2001 /00587 /07  
P.O. Box 3131 Kenmare 1745

Reg. No./Nr.: L 9566 Act/Wet No 36 of/van 1947  
**A READY-FOR-USE  
RODENTICIDE PACKED IN "T" BAGS  
FOR THE CONTROL OF RATS AND  
MICE INDOORS AND OUTDOORS**



**WARNINGS:**

Handle with care. Prevent access to bait by children and domestic animals. Store under lock and key away from food and feedstuffs. Poisonous if swallowed and upon prolonged contact with skin. Toxic to fish and wildlife. In case of poisoning, call a doctor and make this label available to him.

**PRECAUTIONS:**

If you feel unwell, seek medical advice (show the label where possible). Wash hands and exposed skin after handling. Store in original container, tightly closed and in a safe place. Avoid contamination of any food, food utensils, feeds and drinking water. Set bait stations where these will be inaccessible to children and domestic animals. Do not eat, drink, or smoke while handling the bags. Destroy empty container and do not re-use it for any other purpose. Ensure that all the bait is consumed before disposing of the empty container. Dispose of dead rodents by picking it up with rubber or plastic gloves, placing the carcasses in a plastic bag containing sufficient general-purpose household disinfectant to thoroughly wet the carcasses. Seal the bag and dispose of in a safe manner.

**SYMPTOMS OF POISONING:**

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

**FIRST AID TREATMENT:**

If swallowed induce vomiting by tickling the back part of the throat. Repeat until vomit is clear. Take patient IMMEDIATELY to nearest physician.

**NOTE TO PHYSICIAN:**

Intravenous Vitamin K1 10-20mg depending on prothrombin time. In severe cases small transfusions of fresh blood.

**DIRECTIONS FOR USE:**


Use only as directed.

Bait bags should be used unopened as the rodents will also eat the bag material. Place bait where there are signs of rodent activity, like fresh droppings, inside or near burrows, runways and rodent feeding places. Place bait points throughout the infested area. Store unused bags in this pack, close securely and keep in a safe place out of reach of children and animals.

- For rat infestations use bait points of up to 10 bags. Place bait points 10 metres apart, reducing to 5 metres in cases of high infestation.
- For mouse infestations use bait points of 2 bags. Place bait points 5 metres apart, reducing to 2 metres in cases of high infestation.

Protect bait points from non-target animals and from water, preferably using commercially available bait stations. Where these are not available, place the bait on bait trays inside pieces of drainage pipe or under sections of slate, board or corrugated iron. Bait bags may be fixed by "spiking" onto wire inside bait stations. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait.

Do not increase the bait point number. Precautions should be taken to prevent domestic animals from having access to the bait. The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. Rodex Paste Bait is effective against strains of rodent resistant to earlier anticoagulants such as warfarin etc.



Date of Manufacture / Vervaardigingsdatum:

Batch No.: / Lot Nr.: